

AD-A186 499

ON THE MAXIMUM NUMBER OF CONSTRAINTS IN ORTHOGONAL  
ARRAYS(U) ILLINOIS UNIV AT CHICAGO CIRCLE STATISTICAL  
LAB A MEDAYAT ET AL. JUL 87 87-3 AFOSR-TR-87-8320

1/1


UNCLASSIFIED

AFOSR-85-8320

F/G 12/3

NL





DTIC FILE COPY

1



# THE UNIVERSITY OF ILLINOIS AT CHICAGO

AFOSR-TR- 87-1090

AD-A186 499

On the Maximum Number of Constraints  
in Orthogonal Arrays\*

BY

A. HEDAYAT AND J. STUFKEN  
UNIVERSITY OF ILLINOIS AT CHICAGO  
AND UNIVERSITY OF GEORGIA

JULY 1987.

STATISTICAL LABORATORY TECHNICAL REPORT  
No. 87-3.

This document has been approved  
for public release and sale; its  
distribution is unlimited.

DTIC  
ELECTE  
OCT 27 1987  
S D  
OE

DEPARTMENT OF MATHEMATICS,  
STATISTICS, AND COMPUTER SCIENCE

1

**On the Maximum Number of Constraints  
in Orthogonal Arrays\***

BY

A. HEDAYAT AND J. STUFKEN  
UNIVERSITY OF ILLINOIS AT CHICAGO  
AND UNIVERSITY OF GEORGIA

JULY 1987.

STATISTICAL LABORATORY TECHNICAL REPORT  
No. 87-3.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



**DTIC  
ELECTE**  
S 057 0 7 1987  
E

\*Research is supported by Grant AFOSR 85-0320.

6c. ADDRESS (City, State and ZIP Code) P.O. Box 4348, Chicago IL. 60680		7b. ADDRESS (City, State and ZIP Code) <i>BIC 411C</i> Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSOR 85-0320	
8c. ADDRESS (City, State and ZIP Code) <i>BIC 411C</i> Bolling AFB DC 20332-6448		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304
		TASK NO. <i>AS</i>	WORK UNIT NO.
11. TITLE (Include Security Classification)			
12. PERSONAL AUTHOR(S) A. Hedayat and J. Stufken			
13a. TYPE OF REPORT Interim	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Yr. Mo., Day) July, 1987	15. PAGE COUNT 5
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	

Fractional factorial, orthogonal arrays, designs for quality control

# On the Maximum Number of Constraints in Orthogonal Arrays\*

BY

A. HEDAYAT AND J. STUFKEN

UNIVERSITY OF ILLINOIS AT CHICAGO

AND

UNIVERSITY OF GEORGIA

## ABSTRACT

In this paper we show that Bush's bound for the maximum number of constraints in an orthogonal array of index unity is uniformly better than Rao's bound. In addition it is shown, using an argument similar to that needed in the proof of the above result, that Noda's characterization of parameters in orthogonal arrays of strength 4 achieving equality in Rao's bound, leads easily to a similar characterization in arrays of strength 5. <sup>1</sup>

---

AMS 1980 *subject classifications*. Primary 62K15; secondary 05B15. *Keywords and phrases*. Fractional factorials, orthogonal arrays.

<sup>1</sup>Research is supported by Grant AFOSR 85-0320.

**1. Introduction.** Orthogonal arrays were introduced by Rao (1947a) because of their statistical properties when used as fractional factorial designs. See also Rao (1946, 1947b). Rao's papers have been a source of inspiration for many other researchers, who continued to study these arrays and various generalizations. It is mainly due to the work by Taguchi and his colleagues that orthogonal arrays have gained a renewed interest in industrial experimentation for product improvement during recent years, also among researchers in this country.

For the sake of completeness we start with a formal definition of an orthogonal array.

**Definition.** A  $k \times N$  array with entries from  $S = \{0, 1, \dots, s-1\}$  is called an orthogonal array of strength  $t$  and index  $\lambda$  if in any  $t$  rows of the array each (ordered)  $t$ -tuple based on  $S$  appears  $\lambda$  times.

We will denote such an array by  $OA(N, k, s, t)$ . Observe that  $N = \lambda s^t$ . Often  $N$  is called the number of runs,  $k$  the number of constraints, and  $s$  the number of levels.

One of the more interesting problems, both from a mathematical and statistical point of view, is the following. For given values of  $N, s$  and  $t$ , what is the maximum number of constraints that can be accommodated in an orthogonal array? Various results on upper bounds for this number are known. Some are for very general settings, others for very specific ones. For some it is known that equality can be achieved, for others this is an open problem. In this paper we will compare two such upper bounds, one by Rao (1947a), the other by Bush (1952). Rao's result may be stated as follows.

**Theorem 1.** *In an  $OA(\lambda s^t, k, s, t)$  the following inequalities hold.*

*If  $t = 2u$ ,*

$$\lambda s^t \geq 1 + \binom{k}{1}(s-1) + \binom{k}{2}(s-1)^2 + \dots + \binom{k}{u}(s-1)^u.$$

*If  $t = 2u + 1$ ,*

$$\lambda s^t \geq 1 + \binom{k}{1}(s-1) + \binom{k}{2}(s-1)^2 + \dots + \binom{k}{u}(s-1)^u + \binom{k-1}{u}(s-1)^{u+1}.$$

The upper bound for  $k$ , implicitly given by these inequalities, is the most general result on the maximum number of constraints in the literature. Improvements have been

obtained for special cases of the parameters  $N, s$  and  $t$ . Bush (1952) considered the special, but still important case of  $N = s^t$ , arrays of index unity. He proved the following result.

**Theorem 2.** *In an  $OA(s^t, k, s, t)$  the following inequalities hold:*

$$\begin{cases} k \leq t + 1 & \text{if } s \leq t, \\ k \leq s + t - 1 & \text{if } s > t \text{ and } s \text{ is even, or if } t = 2, \\ k \leq s + t - 2 & \text{if } s > t \geq 3 \text{ and } s \text{ is odd.} \end{cases}$$

Improvements on this result for special cases have been obtained, for example by Kounias and Petros (1975). Other improvements result from studying the case  $t = 2$ , a problem that leads its own life. We refer the reader to Dénes and Keedwell (1974) for more on this.

Numerical examples seem to indicate that for orthogonal arrays of index unity the bound for the maximum number of constraints from Theorem 2 is sharper than that from Theorem 1 (Bush (1952)). Personal communications (Bush (1986) and Raghavarao (1986)) made the authors aware that there is apparently no analytical proof of this observation. The main objective of this paper is to provide such a proof, and use the technique in the proof to extend a result by Noda (1979).

**2. Main results.** The following result provides a comparison of the bounds in Theorems 1 and 2.

**Theorem 3.** *For orthogonal arrays of index unity, Bush's bound on the maximum number of constraints is uniformly better than Rao's bound.*

**Proof.** It is well known and easy to show that if  $s \leq t$  the maximum number of constraints equals  $t + 1$ . So we can restrict ourselves to the case  $s > t$ . To prove the theorem it suffices to show that if we use  $k = s + t - 1$  in the inequalities of Theorem 1, we obtain valid inequalities. We distinguish between  $t = 2u$  and  $t = 2u + 1$ .

**Case 1.**  $t = 2u$ . We like to show that  $s^{2u} \geq 1 + \binom{s+2u-1}{1}(s-1) + \dots + \binom{s+2u-1}{u}(s-1)^u$ . First notice that

$$\binom{u}{i}(s+1)^i \geq \binom{s+2u-1}{i}, \quad i = 0, 1, \dots, u.$$



This is obvious for  $i = 0$ . For  $1 \leq i \leq u-1$  it follows if we can show that

$$(u-j)(s+1) \geq s+2u-j-1, \quad j = 0, \dots, u-2,$$

or

$$(u-j-1)s \geq u-1, \quad j = 0, \dots, u-2.$$

This is true since  $j \leq u-2$  and  $s > 2u$ . Finally, for  $i = u$  the above inequality follows since

$$\begin{aligned} u!(s+1)^u &= u(s+1) \cdot (u-1)(s+1) \dots 2(s+1) \cdot (s+1) \geq \\ &u(s+1) \cdot (s+2u-2) \dots (s+u+1)(s+1) \geq \\ &(s+2u-1)(s+2u-2) \dots (s+u+1)(s+u). \end{aligned}$$

The proof for Case 1 is now completed as follows.

$$\begin{aligned} s^{2u} &= (s^2 - 1 + 1)^u = \sum_{i=0}^u \binom{u}{i} (s^2 - 1)^i = \sum_{i=0}^u \binom{u}{i} (s+1)^i (s-1)^i \\ &\geq \sum_{i=0}^u \binom{s+2u-1}{i} (s-1)^i. \end{aligned}$$

**Case 2.**  $t = 2u + 1$ . Here we like to show that

$$s^{2u+1} \geq 1 + \binom{s+2u}{1} (s-1) + \dots + \binom{s+2u}{u} (s-1)^u + \binom{s+2u-1}{u} (s-1)^{u+1}.$$

As in Case 1 we have that  $(s+1)^u \geq \binom{s+2u-1}{u}$ . Observe further that

$$\binom{u}{i-1} (s+1)^{i-1} + \binom{u}{i} (s+1)^i \geq \binom{s+2u-1}{i-1} + \binom{s+2u-1}{i} = \binom{s+2u}{i}.$$

Hence,

$$\begin{aligned} s^{2u+1} &= s \sum_{i=0}^u \binom{u}{i} (s+1)^i (s-1)^i \\ &= \sum_{i=0}^u \binom{u}{i} (s+1)^i (s-1)^{i+1} + \sum_{i=0}^u \binom{u}{i} (s+1)^i (s-1)^i \\ &= 1 + \sum_{i=1}^u \left[ \binom{u}{i-1} (s+1)^{i-1} + \binom{u}{i} (s+1)^i \right] (s-1)^i + (s+1)^u (s-1)^{u+1} \\ &\geq 1 + \sum_{i=1}^u \binom{s+2u}{i} (s-1)^i + \binom{s+2u-1}{u} (s-1)^{u+1}. \end{aligned}$$

This concludes the proof of Case 2 and establishes the result.  $\square$

Similar as in the proof for Case 2 it can be shown that

$$s[1 + \binom{k}{1}(s-1) + \dots + \binom{k}{u}(s-1)^u] = \sum_{i=0}^u \binom{k+1}{i}(s-1)^i + \binom{k}{u}(s-1)^{u+1}.$$

Hence

$$\lambda s^{2u} \geq \sum_{i=0}^u \binom{k}{i}(s-1)^i$$

if and only if

$$\lambda s^{2u+1} \geq \sum_{i=0}^u \binom{k+1}{i}(s-1)^i + \binom{k}{u}(s-1)^{u+1}.$$

Equality can only occur simultaneously. Together with the simple fact that the existence of an  $OA(N, k, s, t)$  implies the existence of an  $OA(N/s, k-1, s, t-1)$ , we are now ready to prove the following result.

**Theorem 4.** *The only values of  $N, k$  and  $s$  for which an  $OA(N, k, s, 5)$  with equality in Rao's inequality can exist, are:*

- (i)  $(N, k, s) = (32, 6, 2)$ , or
- (ii)  $(N, k, s) = (729, 12, 3)$ , or
- (iii)  $(N, k, s) = (27a^2(9a^2 - 1), \frac{3}{5}(3a^2 + 2), 6)$ , where  $a \equiv 0 \pmod{3}$ ,  $a \equiv \pm 1 \pmod{5}$  and  $a \equiv 5 \pmod{16}$ .

**Proof.** An  $OA(N, k, s, 5)$  implies the existence of an  $OA(N/s, k-1, s, 4)$ . By our previous observation, equality in Rao's inequality for the first array occurs if and only if it occurs for the second. The result follows now by the characterization of those cases where equality is possible for arrays of strength 4 by Noda (1979).  $\square$

The arrays corresponding to (i) and (ii) in Theorem 4 are known to exist, see for example Gulati (1971). Finally we point out that an argument as in the proof of Theorem 4 fails for the case  $t = 6$ .

## References

Bush, K.A. (1952). Orthogonal arrays of index unity. *Ann. Math. Statist.* **23** 426-434.

Bush, K.A. (1986). Personal communication.

Dénes, J. and Keedwell, A.D. (1974). *Latin Squares and Their Applications*. Academic Press, New York-London.

Gulati, B.R. (1971). Orthogonal arrays of strength five. *Trabajo Estadíst.* **22** 51-77.

Kounias, S. and Petros, C.I. (1975). Orthogonal arrays of strength three and four with index unity. *Sankhyā Ser. B* **37** 228-240.

Noda, R. (1979). On orthogonal arrays of strength 4 achieving Rao's bound. *J. London Math. Soc. (2)* **19** 385-390.

Raghavarao, D. (1986). Personal communication.

Rao, C.R. (1946). Hypercubes of strength 'd' leading to confounded designs in factorial experiments. *Bull. Calc. Math. Soc.* **38** 67-78.

Rao, C.R. (1947a). On a class of arrangements. *Proc. Edinburgh Math. Soc. (2)* **8** 119-125.

Rao, C.R. (1947b). Factorial experiments derivable from combinatorial arrangements of arrays. *Suppl. J. Roy. Statist. Soc.* **9** 128-139.

END

FEB.

1988

DTIC